EFFECT OF BAY LEAF ETHANOL EXTRACT ON BLOOD GLUCOSE LEVEL IN PATIENTS WITH TYPE 2 DIABETES MELLITUS

Tri Widyawati¹⁾, M. Aron Pase²⁾, Milahayati Daulay³⁾,
Imam Bagus Sumantri⁴⁾

¹⁾Department of Pharmacology and Therapeutic,
Faculty of Medicine, Universitas Sumatera Utara

²⁾Department of Internal Medicine, Faculty of Medicine, Universitas Sumatera Utara

³⁾Department of Physiology, Faculty of Medicine, Universitas Sumatera Utara

⁴⁾Department of Pharmacy Biology, Faculty of Pharmacy, Universitas Sumatera Utara

ABSTRACT

Background: Previous studies showed that bay leaf or *Syzygium polyanthum* (Wight) Walp ethanol extract (SPEE) had antioxidant activity. It is safe for use in human. This study aimed to investigate SPEE effect on fasting blood glucose level (FBGL) among patients with type 2 diabetes mellitus.

Subjects and Method: This was a pilot study using a randomized controlled trial (RCT) design. A total sample of 8 patients with type 2 diabetes mellitus were selected for this study. The intervention group were given 350 mg SPEE in capsule, once daily for 14 days. The control group were given placebo (Pl). The dependent variable was fasting blood glucose level (FBGL). The data were collected at day 0 and after 14 days administration. The effect of EESP was determined by comparing mean FBGL in the EESP-group and Pl-group after intervention. This effect is also shown by Effect Size (ES). The effect is large if $ES > 0.3$. The mean difference after intervention was tested by Mann Whitney test. The means of FBGL of the two groups should be comparable as a result of randomization.

Results: After the intervention, the FBGL (mg/dL) was lower in the EESP-group (Mean= 169.75; SD= 43.72) than the Pl-group (Mean= 225.25; SD= 73.48), and it was not statistically significant ($p= 0.14$). This statistical non-significant effect was anticipated in the pilot study, due to the very small sample size ($n=8$). Effect Size= -0.91, indicating large effect of EESP in lowering the fasting blood glucose level (large if Effect Size > 0.30).

Conclusion: *Syzygium polyanthum* (Wight) Walp ethanol extract (EESP) is effective to lower fasting blood glucose level in patients with type 2 diabetes mellitus.

Keywords: *Syzygium polyanthum*, bay leaf, blood glucose level, diabetes mellitus

Correspondence:

Tri Widyawati. Department of Pharmacology and Therapeutic, Faculty of Medicine, Universitas Sumatera Utara, Medan, 20155, Indonesia. Email: tri.widyawati@usu.ac.id/tw_rozan@yahoo.com. Mobile: 08126582932.

BACKGROUND

Diabetes mellitus (DM) is one of metabolic diseases that characterized by the high level of blood glucose level (BGL) (American Diabetes Association, 2018). This condition affects the human body at multiple organ levels (Bharti et al., 2018). The treatment has become more challenging as conventional oral antidiabetic agents are heterogeneous in their mode of action and causes unwanted effects such as hypoglycaemia, abdominal

discomfort, lactic acidosis, etc. (Ghadge and Kuvalekar, 2017). Long term treatment was also as an important issue that related to the patient adherence (Awodele and Osuolale, 2015; Polonsky and Henry, 2016).

Many plants are traditionally believed to heal diabetes (Bharti et al., 2018), one of which is Salam leaf (bay leaf). Salam leaf is a plant that easily found in Indonesia. Its leaves commonly used as cuisine. This plant has Latin name as *Syzygium polyanthum*

(Wight.) (*S. polyanthum*), which synonyms as *Eugenia balsamea* Ridley, *Eugenia nitida* Duthie and *Eugenia polyantha* Wight (Widyawati et al., 2015a). Other local names are ubar serai, meselengan (Sumatera); samak, kelat samak, dan manting (Jawa) and serah (Malaysia) (Harismah, 2017; Har and Intan, 2012). The efficacy of this plant was known related to its active compounds ie eugenol, *methyl chavicol*, flavonoids, alkaloids, tannin, steroids, triterpenoid, and squalene (Hidayati et al., 2017; Widyawati et al., 2015b; Silalahi, 2017).

Utilization of medicinal plants in the community needs to be supported by its safety data. Pre-clinical study showed that *S. polyanthum* leaf infusion for 28 days was not affecting ureum and creatinine level of Swiss webster mice. At dose 4.5g dried leaves safe to be consumed by human (Kurniawati and Sianturi, 2016). Antihyperglykemic activity of ethanolic extract *S. polyanthum* leaf (30% and 70%) in glucose loading male rabbits showed significant effect (Wahyono, 2008).

Widyawati et al. (2015a) demonstrated that *S. polyanthum* methanol extract 1 g/kg had antihyperglycemic effect in streptozotocin-induced diabetic rats. Studies of its mechanism of actions show the inhibitory activity of the enzyme dipeptidyl Peptidase-IV, alpha glycosidase and alpha amylase and inhibiting glucose absorption in jejunum (Widyawati et al., 2015b; Permatasari and Yuniarni, 2015).

Our previous study showed that 28 days administration of 350 mg of *S. polyanthum* ethanol extract safe to be used in healthy human volunteer. Therefore, the present study was conducted to evaluate its activity in type 2 diabetic patients.

SUBJECT AND METHODS

1. Study Design

This was a pilot study using a randomized controlled trial (RCT) design. The study was

carried out at Medical Faculty, Pharmacy Faculty-Universitas Sumatera Utara and Puskesmas Glugur Darat, Medan, Indonesia from April-September 2019.

2. Population and Sample

A total of 8 patients with type 2 diabetes mellitus were selected for this study using inclusion and exclusion criteria. Inclusion criteria i.e.: diabetic type 2 patients; fasting BGL >126 mg/dl; do not smoke, not taking drugs that may affect BGL in acceptable range and signed the informed consent. The patients that showed any adverse effect or subjective symptoms that cannot be tolerated were excluded from the study. The patients were randomly divided to get either SPEE or Placebo (Pl).

3. Study Method

S. polyanthum fresh leaves were obtained from Kedai Durian, Medan, Indonesia. The plant was identified by Herbarium Medanense (MEDA), Universitas Sumatera Utara. The fresh leaves were washed with running water followed by drying under shade (30-40°C). The dried leaves then were grinded to form powder. To obtain extract (SPEE), the powder were extract using ethanol 96%. The extract then were formulated in capsule at dose 350 mg.

The treatment was administered once daily for 14 days accompanied by oral antihyperglycemic agent, metformin 500 mg twice daily. Blood glucose levels were measured at day 0 (before intervention) and after 14 days of taking capsules. During the administration of the treatment, any complaints experienced was recorded every day. Data were analysed using Mann-Whitney and Wilcoxon sign rank test.

RESULTS

During administration all volunteer completed the study. In EESP-treated group, there were no complaints during observation, while in Pl-treated group, at day 1 as follows

confuse (1 subject (A8), sleepy (1 subject (A11), numb (1 subject (A20). Afterwards A8 mentioned itchy, polyuria, polyphagia.

As shown in Table 1 and Figure 1-2, after the intervention, the FBGL (mg/dL) was lower in the EESP-group (169.75 ± 43.72) than the PL-group (225.25 ± 73.48). FBGL of EESP-treated group decreased (8.85%; BGL difference 16.5 mg/dL) from 186.25 ± 58.57 to 169.75 ± 43.72 , contrarily PL-treated group increased (11.23%; BGL difference 22.75

mg/dL) from 202.5 ± 50.17 to 225.25 ± 73.48 .

Statistically, there was no significant different between PL- and EESP-treated group at the end of the study ($p = 0.14$). This statistical non-significant effect was anticipated in the pilot study, due to the very small sample size ($n=8$). We calculated the effect Size = -0.91 that indicated large effect of EESP in decreasing the FBGL (large if Effect Size > 0.30).

Table 1. Effect of EESP on Fasting Blood Glucose Level

Group	FBGL (mg/dL)		p^b
	Day 0	Day 15	
PL	202.5 ± 50.17	225.25 ± 73.48	0.27
EESP	186.25 ± 58.57	169.75 ± 43.72	0.46
p^a	0.76	0.14	

(a: Kruskal-Wallis; b: Wilcoxon sign rank test)

Figure 1. Blood glucose level difference

Figure 2. Percentage of blood glucose level difference

DISCUSSION

Fasting blood glucose level is one of parameters to diagnose type 2 diabetes mellitus (American Diabetes Association, 2018) which the value should below 126 mg/dL (Wilson and Sexton, 2003). In the present study showed that the diabetic volunteers had high BGL to confirm their diabetic uncontrolled condition. After EESP 14 days administration, the FBGL decreased to 8.85 % or achieved to 16.5 mg/dL reduction.

Suharmiati and Roosihermiatie (2012) reported that metformin in combination to the 700 mg extract of *Andrographis paniculata* and *Syzygium polyanthum* mixture (1:1) for 28 days could significantly decrease FBGL in comparison to control group given metformin with placebo. The FBGL difference was around 37.3 mg/dL.

The higher effect in lowering FBGL of their study probably due to the long term administration. The present study suggested that EESP alone was comparable to Suharmiati and Roosihermiatie study due to the dose and duration of treatment. However, identifying effective drug combinations that significantly improve over single agents still a challenging problem (van Griensven et al., 2010). More drug used more cost spent.

The decreasing of FBGL in the present study insignificant statistically due to the

number of sample as it is for the pilot study. The necessary sample size to achieve stable estimates for correlations depends on some parameters, one of which is the effect size (Schönbrodt and Perugini, 2013). The effect size calculation showed that the EESP has potency as antihyperglycemic agent. This study concluded that *Syzygium polyanthum* (Wight.) Walp ethanol extract (SPEE) is effective to lower fasting blood glucose level in patients with type 2 diabetes mellitus.

ACKNOWLEDGEMENT

The study was supported by Directorate of Research and Community Service Directorate General of Research and Development Strengthening Ministry of Research, Technology and Higher Education (No: 24/UN5.-2.3.1/PPM/KP-DRPM/2019).

REFERENCES

- American Diabetes Association. (2018). Glycaemic targets: standards of medical care in diabetes—2018. *Diabetes Care*, 41 (1): S55-S64.
- Awodele O, Osulale JA (2015). Medication adherence in type 2 diabetes patients: study of patients in Alimosho General Hospital, Igando, Lagos, Nigeria. *African health sciences*, 15: 513-522.
- Bharti SK, Krishnan S, Kumar A, Kumar A.

- (2018). Antidiabetic phytoconstituents and their mode of action on metabolic pathways. *Therapeutic Advances in Endocrinology and Metabolism*, 9: 81-100.
- Ghadge AA, Kuvalekar AA (2017). Controversy of oral hypoglycemic agents in type 2 diabetes mellitus: Novel move towards combination therapies. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 11: S5-S13.
- Harismah K (2017). Pemanfaatan daun salam (*Eugenia polyantha*) sebagai obat herbal dan rempah penyedap makanan. *Warta Lpm*, 19: 110-118.
- Har L, Intan SI (2012). Antioxidant activity, total phenolics and total flavonoids of *Syzygium polyanthum* (Wight) Walp leaves. *International Journal of Medicinal and Aromatic Plants*, 2: 219-228.
- Hidayati MD, Ersam T, Shimizu K, Fatmawati S (2017). Antioxidant activity of *Syzygium polyanthum* extracts. *Indonesian Journal of Chemistry*, 17: 49-53.
- Kurniawati E, Sianturi CY (2016). Manfaat Sarang Semut (*Myrmecodia pendans*) sebagai Terapi Anti diabetes. *Jurnal Majority*, 5:38-42.
- Permatasari D, Yuniarni U (2015). Uji efektifitas antihiperurisemia ekstrak etanol daun salam dan daun jambang serta kombinasinya pada tikus wistar jantan. *Prosiding KNMSA 2015*.
- Polonsky WH, Henry RR (2016). Poor medication adherence in type 2 diabetes: recognizing the scope of the problem and its key contributors. *Patient preference and adherence*, 10, 1299.
- Schönbrodt FD, Perugini M (2013). At what sample size do correlations stabilize? *Journal of Research in Personality*, 47: 609-612.
- Silalahi M (2017). *Syzygium polyanthum* (Wight) Walp. *Jurnal Dinamika Pendidikan*, 10: 1-16.
- Suharmiati S, Roosierhermatie B. (2012). Studi pemanfaatan dan keamanan kombinasi metformin dengan ekstrak campuran *Andrographis paniculate* dan *Syzygium polyanthum* untuk pengobatan diabetes mellitus (Preliminary Study). *Buletin Penelitian Sistem Kesehatan*, 15.
- Wahyono S (2008). Aktivitas hipoglikemia ekstrak etanolik daun salam (*Syzygium polyanthum* (Wight).Walp) dan pengaruhnya terhadap stimulasi parasimpetik pada kelinci jantan yang dibebani glukosa.
- Widyawati T, Purnawan WW, Atangwho IJ, Yusoff NA, Ahmad M, Asmawi MZ (2015a). Anti-diabetic activity of *Syzygium polyanthum* (Wight) leaf extract, the most commonly used herb among diabetic patients in Medan, North Sumatera, Indonesia. *Int. J. Pharm. Sci. Res*, 6, 1698.
- Widyawati T, Yusoff N, Asmawi M, Ahmad M (2015b). Antihyperglycemic effect of methanol extract of *Syzygium polyanthum* (Wight.) leaf in streptozotocin-induced diabetic rats. *Nutrients*, 7: 7764-7780.
- Wilson SJ, Sexton DJ (2003). Elevated preoperative fasting serum glucose levels increase the risk of postoperative mediastinitis in patients undergoing open heart surgery. *Infection Control & Hospital Epidemiology*, 24: 776-778.
- van Griensven, J, Balasegaram M, Meheus F, Alvar J, Lynen L, Boelaert M. (2010). Combination therapy for visceral leishmaniasis. *The Lancet Infectious Diseases*, 10: 184-194.